

POPULATION SYNTHESIS OF NEUTRON STARS, STRANGE (QUARK) STARS AND BLACK HOLES

K. Belczynski^{1,2}, T. Bulik¹, and W. Kluźniak¹

¹Nicolaus Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland

²Harvard-Smithsonian Center for Astrophysics, 60 Garden St., MS. 51, Cambridge, MA 02138, USA

Abstract

We compute and present the distribution in mass of single and binary neutron stars, strange stars, and black holes. The calculations were performed using a stellar population synthesis code. We follow all phases of single and binary evolution, starting from a ZAMS binary and ending in the creation of one compact object (neutron star, black hole, strange star) and a white dwarf, or two compact objects (single or binary). We assume that neutron stars are formed in the collapse of iron/nickel cores in the mass range $M_0 < M < M_1$, quark stars in the range $M_1 < M < M_2$, and black holes for core masses $M > M_2$ and find that the population of quark stars can easily be as large as the population of black holes, even if there is only a small mass window for their formation.

Key words: compact objects – masses, binaries – population synthesis.

1. INTRODUCTION

Binary population synthesis is a useful tool for studying the statistical properties of stars, including the compact objects. In this paper we wish to address the following questions: what is the distribution of masses of the compact objects? what is the relation between the stellar initial mass and the final mass of a compact object, when binary evolution is taken into account? given the distribution of compact object masses, what are the relative numbers of different types of objects (neutron stars, quark stars, black holes)?

In section 2 we shortly describe the population synthesis code used here, in section 2 we present the results, and finally in section 4 we present the conclusions.

2. POPULATION SYNTHESIS CODE

We use the stellar binary population synthesis code consisting of two parts. The single star evolution is based on modified formulae from Hurley et al. (2000). We have changed the prescription for mass of the compact object formed in a supernova explosion. We use original Hurley et al. (2000) formulae to obtain final CO core mass. We use models of Woosley (1986) to calculate the final FeNi core mass (for a given CO core mass), which will collapse and form a compact object during supernova explosion. Finally, we include calculations of Fryer and Kalogera (2000) to take into account black hole formation both through fall back and directly.

The binary evolution is described in Belczyński et al. (2000). We evolve only binaries where at least one star will undergo a supernova explosion. The evolution starts at ZAMS. During the course of evolution we include the following effects:

- wind mass loss (standard, Wolf-Rayet, LBV)
- tidal circularization of binary orbit
- magnetic breaking
- conservative/nonconservative mass transfer
- common envelope evolution
- rejuvenation
- hyper-accretion onto compact objects
- detailed supernova explosion (SN) treatment
- partial and complete fall back onto compact objects in SN

Many binaries are disrupted in supernova explosions, as a result of mass loss and the natal kick. For supernova kicks we use distribution presented by Cordes and Chernoff (1997). We continue to evolve all stars, both single and binary components, until the formation of a stellar remnant.




Figure 1. The various routes of compact object formation (compact object = a neutron star, a strange star or a black hole).

3. RESULTS

In Figure 1 we show different formation routes of compact objects: neutron stars, strange stars and black holes. Compact objects are formed both through single and binary stellar evolution. Single compact objects are descendants of massive single stars but may also be formed as a binary system is disrupted in a supernova explosion of one of its components (these are the paths on the left hand side of Fig.1). We will denote the single compact objects formed as a result of the binary evolution as group I. Under favorable conditions some binaries survive supernova explosions, and they finally form tight systems with compact object/objects. Most of these binaries will consist of a white dwarf and a compact object and the rest will form binaries with two compact objects (we will denote the compact object in binaries with white dwarfs as group II, and the double compact objects as group III).




Figure 2. The number per mass interval of neutron stars / strange stars / black holes formed along each route shown in Fig.1. The solid line corresponds to the case of single stellar evolution, the dotted line represents the single compact objects formed in binary evolution (group I), the short dashed line shows the compact objects in white dwarf binaries (group II), and the long dashed line - the compact objects in double compact object binaries (group III).

In Figure 2 and 3 we show production rates of compact objects as a function of their final mass for the four different formation routes presented in Fig.1.

Figure 2 shows the number of compact objects per mass interval formed along each route of Figure 1. We start forming compact objects at mass $\sim 1.2M_{\odot}$ and their number falls off with the mass of a final compact object, as expected for our assumed initial mass function $\sim M^{-2.7}$. A peak around $\sim 10M_{\odot}$ reflects our relation between ZAMS mass of a progenitor and the final mass of a compact object, which is shown for single stars with a solid line in Fig.4. This relation for wide range of progenitor ZAMS masses results in a final compact object mass of $\sim 10M_{\odot}$. This is an effect of a stellar wind which increases with a stellar mass, and thus decreases the final mass of a compact object flattening out the relation at about $10M_{\odot}$.




Figure 3. The cumulative fraction of compact objects (neutron stars, strange stars, black holes) corresponding to the differential distribution of Fig.2. The line types are the same as in Figure 2.

Figure 3 shows cumulative fraction of compact objects as a function of their final mass for different formation routes. Note the very fast initial rise of cumulative curves for single compact remnants and remnants white dwarf binaries. Most of compact objects in these groups have mass below $2M_{\odot}$. Only for double compact objects a large fraction of remnants have higher final masses.

In Figure 4 we plotted the mass of a single compact object as a function of the ZAMS mass of a progenitor star. Single compact objects coming from single stars are shown with solid line, and the ones coming from disrupted binary systems are shown with points. Points which are over the solid line form a group of compact objects whose progenitors gained mass in binary interactions, and thus they formed more massive compact objects then corresponding single stars of the same ZAMS mass. Points which are below the solid line represent a group of compact objects whose progenitors were donors in different mass transfer events, and thus they lost part of their mass and formed less massive compact objects. There is also a group of compact objects com-

ing from binary progenitors, which follows exactly $M_{\text{zams}}-M_{\text{final}}$ relation for compact objects formed from single stars. This group comes from wide, non interacting binaries, which in the most cases were disrupted in a first supernova explosion. The exploding component was unaffected by its companion so it has left behind a compact object remnant exactly as it would be formed from a single star. Then its companion, also unaffected by binary evolution, continues its evolution. Provided that it is massive enough, it goes through a supernova explosion leaving second single compact object remnant, of a mass fitting the single star $M_{\text{zams}}-M_{\text{final}}$ relation.

4. CONCLUSIONS

We have shown the effects of the binary evolution on the distribution of masses of compact objects. As expected the bulk of the population of compact objects have masses below $2M_{\odot}$. While for single stellar evolution there exists a unique relation between the stellar mass and the compact object mass, there is no




Figure 4. The mass of a single compact object as a function of the ZAMS mass of the progenitor star. The solid line shows the correspondence between the initial and final mass for initially single stars. The points correspond to compact objects whose progenitors were initially in binary systems. The three panels correspond to group I, II, and III, from top to bottom respectively. Note that binary stars can have more than one fate.

such relation when the binary evolution is taken into account. The binary evolution works both ways, the masses of the compact objects formed in binaries can be both smaller and larger than in the corresponding case of single stellar evolution.

The masses of the compact objects are concentrated in the low mass range - see Figure 3. Thus even if the mass interval (M_1, M_2) for the formation of strange (quark) stars is small, the fraction of strange stars in the whole compact object population can be huge. The exact fraction as a function of M_1 and M_2 can be easily read off Figure 3. On the other hand the fraction of black holes hardly depends on M_2 (the cumulative curves flatten above $3 M_\odot$). We conclude that the population of quark stars can easily be as large as the population of black holes, even if there is only a small mass window for their formation.

Acknowledgments

The authors gratefully acknowledge support of the KBN grant 2P03D00418.

References

- Hurley J.R., Pols O.R., Tout C.A., 2000, MNRAS 315, 543
- Belczynski K., Kalogera V., Bulik T., 2000, in preparation
- Cordes J., Chernoff D. F., 1997, ApJ, 482, 971
- Woosley S.E., 1986, in 'Nucleosynthesis and Chemical Evolution', 16th Saas-Fee Course, eds. B. Hauck et al., Geneva Observatory, p.1
- Fryer C.L., Kalogera V., 2000, astro-ph/9911312